

Stage 1 Obstetrical Hemorrhage

Cumulative blood loss 500- 1000cc AND HR >110, BP <85/45, O2 sat <95 – OR- Increased bleeding during recovery or postpartum period

Primary Nurse

Communication	Action
<ul style="list-style-type: none"> ○ Primary/Covering Physician (Provider) ○ Charge Nurse ○ Anesthesia 	<ul style="list-style-type: none"> ○ Obtain primary IV access if not present ○ Vital Signs every 10 minutes ○ Apply Fundal Massage ○ Empty Bladder

Charge Nurse

Communication	Action
<ul style="list-style-type: none"> ○ Primary/Covering Physician(Provider) ○ Anesthesia ○ OR Team 	

Physician/Midwife

Communication	Action
<ul style="list-style-type: none"> ○ Anesthesia ○ Charge Nurse 	<ul style="list-style-type: none"> ○ Evaluate Patient ○ Order uterotonics (Pitocin, Methergine, Hemabate, Misoprostol) ○ Order Type and Screen ○ Clear Clot from Uterus and Evaluate for Cervical Lacerations and Retained Placenta

Anesthesia

Communication	Action
	<ul style="list-style-type: none"> ○ Evaluate Patient when Notified ○ Prepare for possibility of additional anesthesia support ○ Coordinate with OB attending